

## 13 Gravitational fields

### 13.1 Gravitational field





Candidates should be able to:

- 1 understand that a gravitational field is an example of a field of force and define gravitational field as force per unit mass
- 2 represent a gravitational field by means of field lines

- When **two or more masses** are in proximity with each other there is an **attractive force** between them.
- This force is called **gravity**
- A **gravitational field** is defined as a **region of space where a mass experience a force due to the gravitational attraction of another mass.**
- The SI unit for gravitational field strength is **N kg<sup>-1</sup>** or **ms<sup>-2</sup>**
- The gravitational field strength ( $g$ ) at a point is the force due to gravity or weight ( $F_g$ ) per unit mass ( $m$ ) of an object at that point:

$$g = \frac{F_g}{m}$$

- The larger the planet, the larger the  $g$ !
- Gravitational field lines like magnetic and electric field lines gives us an indication to their direction.
- Unlike the other two however, gravitational field lines are always attractive and never repulsive!
- Below are some examples of gravitational field lines




## 13.2 Gravitational force between point masses

Candidates should be able to:


- 1 understand that, for a point outside a uniform sphere, the mass of the sphere may be considered to be a point mass at its centre
- 2 recall and use Newton's law of gravitation  $F = Gm_1m_2/r^2$  for the force between two point masses
- 3 analyse circular orbits in gravitational fields by relating the gravitational force to the centripetal acceleration it causes
- 4 understand that a satellite in a geostationary orbit remains at the same point above the Earth's surface, with an orbital period of 24 hours, orbiting from west to east, directly above the Equator

- For a point outside a uniform sphere, the mass of the sphere may be considered to be a point mass at its centre.
- A uniform sphere is one where its mass is **distributed evenly**.
- The gravitational field lines around a uniform sphere are therefore **identical to those around a point mass**
- An object can be regarded as point mass **when a body covers a very large distance as compared to its size**.
- Radial fields are considered **non-uniform** fields.
- Hence  $g$  is different depending on how far you are from the centre of mass of the sphere



- Newton's Law of Gravitation states that the **gravitational force between two point masses is proportional to the product of the masses and inversely proportional to the square of their separation.**
- This can be written as

$$F_G = \frac{Gm_1m_2}{r^2}$$




- Here  $G$  is Newton's gravitational constant  $6.67 \times 10^{-11} \text{ Nm}^2\text{kg}^{-2}$
- This value will be given in exam.
- In order for a planet to stay in orbit around the sun (as oppose to falling into the sun!) the planet must travel in a circular orbit around the sun in order for **centripetal force** to balance the **gravitational force**

$$F_C = F_G$$

$$\frac{m_2 v_t^2}{r} = \frac{Gm_1 m_2}{r^2}$$

$$v_t^2 = \frac{Gm_1}{r}$$

- The equation above proves that all planets travel at same tangential speed ( $v_t$ ) around the sun since the speed is only dependent on the mass of the sun ( $m_1$ )



- Most satellites orbiting the earth follow a **geostationary orbit**.
- The criteria for geostationary orbit are:
  - Remains directly above the equator
  - Moves from west to east (same direction as the Earth spins)
  - Has an orbital time period equal to Earth's rotational period of 24 hours

### 13.3 Gravitational field of a point mass

Candidates should be able to:

- 1 derive, from Newton's law of gravitation and the definition of gravitational field, the equation  $g = GM/r^2$  for the gravitational field strength due to a point mass
- 2 recall and use  $g = GM/r^2$
- 3 understand why  $g$  is approximately constant for small changes in height near the Earth's surface

- In A levels the candidate must be able to derive the gravitational field equation

$$g = \frac{Gm_1}{r^2}$$


- To derive the equation above first take Newton's law of gravitation force equation

$$F_G = \frac{Gm_1m_2}{r^2}$$

And substitute  $F_G$  with

$$F_G = m_2g$$

- $g$  here is the same as acceleration due to gravity that you have been using!
- The SI unit is in  $\text{N kg}^{-1}$  or  $\text{ms}^{-2}$
- Based on the gravitational field equation,  $g$  is directly proportional to the mass of the planet ( $m_1$ ) and inversely proportional to the square of the radius of the planet  $r^2$



- The value of  $g$  changes very little for small changes in height near the surface of the Earth ( $9.81 \text{ ms}^{-2}$ ).
- This is because any height change is very small compared to the radius of the earth ( $r$ )

### 13.4 Gravitational potential

Candidates should be able to:

- 1 define gravitational potential at a point as the work done per unit mass in bringing a small test mass from infinity to the point
- 2 use  $\phi = -GM/r$  for the gravitational potential in the field due to a point mass
- 3 understand how the concept of gravitational potential leads to the gravitational potential energy of two point masses and use  $E_p = -GMm/r$

- Recall that gravitational potential energy (GPE) is given by

$$\text{GPE} = mgh$$

- It is defined as the energy an object possess due to its position in a gravitational field

- We can replace the **height (h)** with the **distance from the center of the earth (r)** and **mass (m)** with the **mass of the object above the earth (m<sub>2</sub>)**

$$GPE = m_2gr$$

- Gravitational potential ( $\phi$ ) is defined **work done per unit mass in bringing a test mass from infinity to a defined point**
- So basically, gravitational potential ( $\phi$ ) is just **GPE per kg<sup>-1</sup> (GPE/mass)!**
- The SI unit is **J kg<sup>-1</sup>**
- Divide the equation above with **m<sub>2</sub>**

$$\phi = gr$$

Recall that

$$g = \frac{Gm_1}{r^2}$$

Substituting into the above equation we get

$$\phi = \frac{-Gm_1}{r}$$

The equation for GPE of two-point masses  $m_1$  and  $m_2$  can thus be written as

$$GPE = \frac{Gm_1m_2}{r}$$

If the object was initially at  $r_1$  from the center of the Earth and then moved further to  $r_2$  Both GPE and  $\phi$  would be

$$\Delta GPE = Gm_1m_2 \left( \frac{1}{r_1} - \frac{1}{r_2} \right)$$

$$\Delta\phi = Gm_1 \left( \frac{1}{r_1} - \frac{1}{r_2} \right)$$