




7 Waves

7.1 Progressive waves


Candidates should be able to:

- 1 describe what is meant by wave motion as illustrated by vibration in ropes, springs and ripple tanks
- 2 understand and use the terms displacement, amplitude, phase difference, period, frequency, wavelength and speed
- 3 understand the use of the time-base and y-gain of a cathode-ray oscilloscope (CRO) to determine frequency and amplitude
- 4 derive, using the definitions of speed, frequency and wavelength, the wave equation $v = f\lambda$
- 5 recall and use $v = f\lambda$
- 6 understand that energy is transferred by a progressive wave
- 7 recall and use $intensity = power/area$ and $intensity \propto (amplitude)^2$ for a progressive wave


- Waves are created when particles oscillate or vibrate.
- **Energy** is transferred through these moving oscillations.



- You can represent waves on **two** types of plots:
 - Displacement-distance plot



- Displacement-time plot




- Below are some terminology for waves:
- Amplitude, A** : The maximum displacement from the original position. The SI unit for amplitude is in meters.
- Wavelength, λ** : The horizontal distance between two **wave fronts**. The SI unit for wavelength is in meters
- Period, T** : Time taken for the wave to complete a cycle or return to its original displacement. The SI unit for periods is seconds.
- Frequency, f** : The number of complete cycle's in a second (How many times did the wave go up, down and up again or down, up, and down again in 1 second). The SI unit for frequency is hertz (Hz) OR seconds⁻¹.


- Hence, relationship between **frequency** and **period** is

$$f = 1/T$$

- A cathode ray oscilloscope (CRO) can be used to display the waveform of electrical signals.



- Below is an example of a CRO display with the x-axis being time and y-axis being voltage



- The period of the wave can be determined from the time based (how

Worked example

Assuming from the diagram above each cm represent 1 second, find the frequency of the wave

- Wave equation is given by

$$v = f\lambda$$

Here v is the speed of the wave, f is the frequency of the wave and λ is the wavelength of the wave

Derivation for the equation above

- Progressive waves transfer **energy**.
- The amount of energy passing per unit time (P) through a unit of area (A) is the intensity, I of the wave

$$I = \frac{P}{A}$$

- The area the wave passes through is perpendicular to the direction of its velocity
- The intensity of a progressive wave is also proportional to its amplitude squared and frequency squared

$$I \propto A^2$$



$$I \propto f^2$$

7.2 Transverse and longitudinal waves



Candidates should be able to:

- 1 compare transverse and longitudinal waves
- 2 analyse and interpret graphical representations of transverse and longitudinal waves

- **Transverse waves** are waves where the particles vibrate **perpendicular** along the lines of motions and consists of a series of "crest" and "troughs".
- Examples include electromagnetic waves, water ripples and vibration on a guitar string.
- **Longitudinal waves** are waves where the particles vibrate **along** the lines of motion and consist of a series of compression and expansions (rarefractions).
- Examples include sound waves.
- Visual and graphical representation of transverse waves



- Visual and graphical representation of longitudinal waves




7.3 Doppler effect for sound waves

Candidates should be able to:

- 1 understand that when a source of sound waves moves relative to a stationary observer, the observed frequency is different from the source frequency (understanding of the Doppler effect for a stationary source and a moving observer is not required)
- 2 use the expression $f_o = f_s v / (v \pm v_s)$ for the observed frequency when a source of sound waves moves relative to a stationary observer

- The sound of a siren changes as it moves closer or further from you.
- This phenomenon is due to the **doppler effect**.
- When the observer and the source sound are both stationary, the waves are at the same frequency for both observer and source.



- Here f' is the observed frequency, f is the frequency from the source, V is the wave velocity and V_s is the source velocity.


7.4 Electromagnetic spectrum

Candidates should be able to:

- 1 state that all electromagnetic waves are transverse waves that travel with the same speed c in free space
- 2 recall the approximate range of wavelengths in free space of the principal regions of the electromagnetic spectrum from radio waves to γ -rays
- 3 recall that wavelengths in the range 400–700 nm in free space are visible to the human eye

- Electromagnetic waves are transverse waves.
- It consists of electric field and magnetic field components.
- It can propagate without the need of a medium to carry them unlike mechanical waves.
- The speed that electromagnetic waves travel at is $3 \times 10^8 \text{ ms}^{-1}$
- If this number seems familiar it's because that's the speed of light. Light is a wave or more specifically an electromagnetic wave.
- There are 7 types of waves in the electromagnetic spectrum.

THE ELECTROMAGNETIC SPECTRUM




7.5 Polarisation

Candidates should be able to:

- 1 understand that polarisation is a phenomenon associated with transverse waves
- 2 recall and use Malus's law ($I = I_0 \cos^2 \theta$) to calculate the intensity of a plane polarised electromagnetic wave after transmission through a polarising filter or a series of polarising filters



- Only transverse waves can be polarized.
- Polarization means that vibrations are restricted to one direction.
- Waves can be polarised through a polariser or polarising filter.
- Example polaroid sunglasses.



- Polarising a wave reduces its amplitude.
- Malus' Law is used to find the intensity of light after passing through a number of filters.

$$I = I_0 \cos^2 \theta$$

Here I is the remaining intensity (W m^{-2}), I_0 is the original intensity, θ is the angle between polarised light and transmission axis (degrees)

Angle of transmission axis θ / degrees	Direction of transmission axis	$\cos^2 \theta$	Transmitted Intensity I / W m^{-2}	Max or min light intensity transmitted
0		1	I_0	Max
180				
90		0	0	Min
270				